


Hybrid approach for iliofemoral obstructive disease



ARISTOTLE
UNIVERSITY



Dr. Nikolaos Melas, PhD

Vascular and Endovascular Surgeon

Military Doctor

Associate in 1st department of Surgery,
Aristotle University of Thessaloniki, Greece

Associate in Interbalcan Medical Center

Previous History


- 72 years old male
- CAD
- DM
- Hypertension
- COPD

Previous history

1998 Fontaine stage IIb (R)
CIA, EIA, IIA occlusion
TASC D



Iliofemoral bypass (R)




Previous history

2005 Fontaine stage IIb bilateral
Proximal anastomosis stenosis (R)
Multifocal (L) CIA, EIA stenosis
Obstruction (L) IIA
TASC C




kissing stent




Current situation


2008 Fontaine stage IIb bilateral
Proximal anastomosis restenosis (R)
Multifocal (L) EIA stenosis
(L) CFA + profunda stenosis
TASC D


- 1) (L) Femoral endarterectomy+Profundoplasty
(Bovine patch)
- 2) SE stent on the (L) EIA + PTA (L) iliac axis
- 3) SE stent on the (R) anastomosis







Initial DSA of current situation





E 0 %
LUT GER_15_L





Alternative solutions

Ilioferm by pass ?

Aorto bifem by pass?